

Hydroxytyrosol Gesunde Effekte

Bibliographie über wissenschaftliche Evidenzen

1. Covas MI, de la Torre R, Fitó M. *Scientific evidence of the benefits of virgin olive oil for human health*. Medicina Balear 2014; 29 (2); 39-46.
2. Granados-Principal S, Quiles JL, Ramirez-Tortosa CL, Sanchez-Rovira P and Ramirez-Tortosa MC. *Hydroxytyrosol: from laboratory investigations to future clinical trials*. Nutrition Reviews Vol. 68(4):191–206
3. Visioli F and Galli C. *Antiatherogenic components of olive oil*. Curr Atheroscler Rep. 2001 Jan;3(1):64-7.
4. Visioli F and Galli C. *Biological properties of olive oil phytochemicals*. Crit Rev Food Sci Nutr. 2002;42(3):209-21.
5. Visioli F, Poli A, Gall C. *Antioxidant and other biological activities of phenols from olives and olive oil*. Med Res Rev. 2002 Jan;22(1):65-75.
6. Sofi F, Cesari F, Abbate R, Gensini GF, Casini A. *Adherence to Mediterranean diet and health status: meta-analysis*. BMJ 2008;337:a1344.
7. Visioli F, Bellomo G, Montedoro G and Galli C. *Low density lipoprotein oxidation is inhibited in vitro by olive oil constituents*. Atherosclerosis. 1995 Sep;117(1):25-32.
8. Deiana M, Aruoma OI, Bianchi ML, Spencer JP, Kaur H, Halliwell B, Aeschbach R, Banni S, Dessi MA and Corongiu FP. *Inhibition of peroxynitrite dependent DNA base modification and tyrosine nitration by the extra virgin olive oil-derived antioxidant hydroxytyrosol*. Free Radic Biol Med. 1999 Mar;26(5-6):762-9.
9. de la Puerta R, Ruiz Gutierrez V and Hoult JR. *Inhibition of leukocyte 5-lipoxygenase by phenolics from virgin olive oil*. Biochem Pharmacol. 1999 Feb 15;57(4):445-9.
10. Kohyama N, Nagata T, Fujimoto S and Sekiya K. *Inhibition of arachidonate lipoxygenase activities by 2-(3,4-dihydroxyphenyl)ethanol, a phenolic compound from olives*. Biosci Biotechnol Biochem. 1997 Feb;61(2):347-50.
11. Petroni A, Blasevich M, Salami M, Papini N, Montedoro GF and Galli C. *Inhibition of platelet aggregation and eicosanoid production by phenolic components of olive oil*. Thromb Res. 1995 Apr 15;78(2):151-60.
12. Bonanome A, Pagnan A, Caruso D, Toia A, Xamin A, Fedeli E, Berra B, Zamburlini A, Ursini F and Galli G., *Evidence of postprandial absorption of olive oil phenols in humans*. Nutr Metab Cardiovasc Dis. 2000 Jun;10(3):111-20.
13. EFSA Journal 2011; 9(4):2033.
14. Xu CL, Sim MK. *Reduction of dihydroxyphenylacetic acid by a novel enzyme in the rat brain*. Biochem Pharmacol. 1995; 50:1333–1337.
15. de la Torre R. *Bioavailability of olive oil phenolic compounds in humans*. Inflammopharmacology. 2008 Oct;16(5):245-7.

16. Edwards DJ, Rizk M. *Conversion of 3, 4-dihydroxyphenylalanine and deuterated 3, 4-dihydroxyphenylalanine to alcoholic metabolites of catecholamines in rat brain.* J Neurochem. 1981;36:1641–1647.
17. Miro-Casas E, Covas MI, Farre M, Fitó M., Ortuno J, Weinbrenner T, Roset P and de la Torre R. *Hydroxytyrosol disposition in humans.* Clin Chem. 2003 Jun;49(6 Pt 1):945-52.
18. Visioli F, Galli C, Grande S, Colonnelli K, Patelli C, Galli G and Caruso D. *Hydroxytyrosol excretion differs between rats and humans and depends on the vehicle of administration.* J Nutr. 2003 Aug;133(8):2612-5.
19. Zainal TA, Weindruch R, Szweda LI and Oberley TD. *Localization of 4-hydroxy-2 nonenal-modified-proteins in kidney following iron overload.* Free Radic Biol Med. 1999 May;26(9-10):1181-93.
20. Vissers MN, Zock PL, Roodenburg AJ, Leenen R, Katan MB. *Olive oil phenols are absorbed in humans.* J Nutr 2002;132:409–17.
21. Bai C, Yan X, Takenaka M, Sekiya K, Nagata T. *Determination of synthetic hydroxytyrosol in rat plasma by GC-MS.* J Agric Food Chem 1998;46:3998–4001.
22. Covas MI, de la Torre K, Farré-Albaladejo M, Kaikkonen J, Fitó M, López-Sabater C, Pujadas-Bastardes MA, Joglar J, Weinbrenner T, Lamuela-Raventós RM, de la Torre R. *Postprandial LDL phenolic content and LDL oxidation are modulated by olive oil phenolic compounds in humans.* Free Radic Biol Med. 2006 Feb 15;40(4):608-16.
23. Corona G., Tzounis X., Dessi M.A., Deiana M., Debnam E.S., Visioli F. and Spencer J.P.E., *The fate of olive oil polyphenols in the gastrointestinal tract: implications of gastric and colonic microfloradependent biotransformation.* Free Radic Res. 2006 Jun;40(6):647-58.
24. Tuck KL, Freeman MP, Hayball PJ, Stretch GL, Stupans I. *The in vivo fate of hydroxytyrosol and tyrosol, antioxidant phenolic constituents of olive oil, after intravenous and oral dosing of labeled compounds to rats.* J Nutr 2001;131:1993–6.
25. Lusis AJ. *Atherosclerosis.* Nature 2000;407:233–41.
26. Bulotta S, Celano M, Lepore SM, Montalcini T, Pujia A, Russo D. *Beneficial effects of the olive oil phenolic components oleuropein and hydroxytyrosol: focus on protection against cardiovascular and metabolic diseases.* J Transl Med. 2014 Aug 3;12:219.
27. Camargo A, Ruano J, Fernandez JM, Parnell LD, Jimenez A, Santos-Gonzalez M, Marin C, Perez-Martinez P, Uceda M, Lopez-Miranda J, Perez-Jimenez F. *Gene expression changes in mononuclear cells in patients with metabolic syndrome after acute intake of phenol-rich virgin olive oil.* BMC Genomics. 2010 Apr 20;11:253.
28. Castañer O, Covas MI, Khymenets O, Nyssonen K, Konstantinidou V, Zunft HF, de la Torre R, Muñoz-Aguayo D, Vila J, Fitó M. *Protection of LDL from oxidation by olive oil polyphenols is associated with a downregulation of CD40-ligand expression*

- and its downstream products in vivo in humans.* Am J Clin Nutr. 2012 May;95(5):1238-44.
29. Cicerale S, Lucas L, Keast R. *Biological activities of phenolic compounds present in virgin olive oil.* Int J Mol Sci. 2010 Feb 2;11(2):458-79.
 30. Samuel SM, Thirunavukkarasu M, Penumathsa SV, Paul D, Maulik N. *Akt/FOXO3a/SIRT1-mediated cardioprotection by n-tyrosol against ischemic stress in rat in vivo model of myocardial infarction: switching gears toward survival and longevity.* J Agric Food Chem. 2008 Oct 22;56(20):9692-8.
 31. Raederslorff D. *Antioxidant activity of olive polyphenols in humans: a review.* Int J Vitam Nutr Res. 2009 May;79(3):152-65
 32. Fabiani R, Rosignoli P, De Bartolomeo A, Fuccelli R, Servili M, Montedoro GF, Morozzi G. *Oxidative DNA damage is prevented by extracts of olive oil, hydroxytyrosol and other olive phenolic compounds in human blood mononuclear cells and HL60 cells.* J Nutr. 2008 Aug;138(8):1411-6.
 33. Pereira-Caro G, Sarriá B, Madrona A, Espartero JL, Goya L, Bravo L, Mateos R. *Alkyl hydroxytyrosyl ethers show protective effects against oxidative stress in HepG2 cells.* J Agric Food Chem. 2011 Jun 8;59(11):5964-76.
 34. Zrelli H, Matsuoka M, Kitazaki S, Zarrouk M, Miyazaki H. *Hydroxytyrosol reduces intracellular reactive oxygen species levels in vascular endothelial cells by upregulating catalase expression through the AMPK-FOXO3a pathway.* Eur J Pharmacol. 2011 Jun 25;660(2-3):275-82
 35. Brunswick Laboratories, 200 Turnpike Road, Southborough, MA 01772, USA.
 36. Haytowitz DB, Bhagwat S. *USDA Database for the Oxygen Radical Absorbance Capacity (ORAC) of Selected Foods, Release 2, May 2010.* U.S. Department of Agriculture - Agricultural Research Service - Beltsville Human Nutrition Research Center - Nutrient Data Laboratory -10300 Baltimore Avenue - Building 005, Room 107, BARC-West - Beltsville, Maryland 20705, USA.
 37. Valls-Pedret C, Lamuela-Raventós RM, Medina-Remón A, Quintana M, Corella D, Pintó X, Martínez-González MÁ, Estruch R, Ros E. *Polyphenol-rich foods in the Mediterranean diet are associated with better cognitive function in elderly subjects at high cardiovascular risk.* J Alzheimers Dis. 2012;29(4):773-82.
 38. Bitler CM, Matt K, Irving M, Hook G, Yusen J, Eagar F, Kirschner K, Walker B, Crea R. *Olive extract supplement decreases pain and improves daily activities in adults with osteoarthritis and decreases plasma homocysteine in those with rheumatoid arthritis.* Nutrition Research 27 (2007) 470-477.
 39. Covas MI, Nyssönen K, Poulsen HE, Kaikkonen J, Zunft HJ, Kiesewetter H, Gaddi A, de la Torre R, Mursu J, Bäuml H, Nascetti S, Salonen JT, Fitó M, Virtanen J, Marrugat J; EUROLIVE Study Group. *The effect of polyphenols in olive oil on heart disease risk factors: a randomized trial.* Ann Intern Med. 2006 Sep 5;145(5):333-41.

40. González-Santiago M et al. *One-month administration of hydroxytyrosol, a phenolic antioxidant present in olive oil, to hyperlipemic rabbits improves blood lipid profile, antioxidant status and reduces atherosclerosis development.* *Atherosclerosis.* 2006 Sep;188(1):35-42.
41. González-Santiago M, Fonollá J, Lopez-Huertas E. *Human absorption of a supplement containing purified hydroxytyrosol, a natural antioxidant from olive oil, and evidence for its transient association with low-density lipoproteins.* *Pharmacol Res.* 2010 Apr;61(4):364-70.
42. Perona JS, Montero E, Sánchez-Domínguez JM, Cañizares J, Garcia M, Ruiz-Gutiérrez V. *Evaluation of the effect of dietary virgin olive oil on blood pressure and lipid composition of serum and low-density lipoprotein in elderly type 2 diabetic subjects.* *J Agric Food Chem.* 2009 Dec 9;57(23):11427-33.
43. Rietjens SJ, Bast A, de Vente J, Haenen GR. *The olive oil antioxidant hydroxytyrosol efficiently protects against the oxidative stress-induced impairment of the NObullet response of isolated rat aorta.* *Am J Physiol Heart Circ Physiol.* 2007 Apr;292(4):H1931-6.
44. Visioli F, Wolfram R, Richard D, Abdullah MI, Crea R. *Olive phenolics increase glutathione levels in healthy volunteers.* *J Agric Food Chem.* 2009 Mar 11;57(5):1793-6.